Feeding Do's & Don'ts



- Hold and cuddle your baby when breastfeeding or feeding a bottle.
- · Smile, talk, and maintain eye contact with your baby at mealtime.
- Make mealtime a relaxed and happy time together.
- Offer a variety of foods including fruits and vegetables at meals and snacks.
- Let baby feed himself, even if it is messy!
- Try foods again if baby "does not like it" at first. It might take 10-15 times!
- Be positive when offering new foods, even if you do not like the food.
- Listen for a "pop" when you open a new jar of baby food. If it doesn't "pop," throw it out.
- Use refrigerated baby food within 3 days.



- No infant feeders. Do not prop the bottle.
- No cereals, juices or foods in the bottle.
- · No baby desserts or mixed dinners.
- No cow's, goat's or soy milk before one year old.
- No fruit drinks, tea, Kool-aid[®], Gatorade™, soda pop or coffee
- Do not feed directly from the baby food jar.
- Do not add salt, sugar, butter, or honey to baby's foods.

Persons with disabilities who require alternative means of communication for should contact the Agency (State or local) who are deaf, hard of hearing or have speech disabilities may contact USDA

To file a program complaint of Discrimination Complaint Form, (AD-3027) found online at: http://www. ascr.usda.gov/complaint_filing_cust.html, and at any USDA office, or write a letter all of the information requested in the form. (866) 632-9992. Submit your completed form or letter to USDA by:

(1) mail: U.S. Department of Agriculture Office of the Assistant Secretary for 1400 Independence Avenue, SW Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

Ohio Department of Health

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, sex, disability, age, or reprisal or retaliation for prior civil rights activity in any program or activity conducted or funded by USDA.

program information (e.g. Braille, large print, audiotape, American Sign Language, etc.), where they applied for benefits. Individuals through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

discrimination, complete the USDA Program addressed to USDA and provide in the letter To request a copy of the complaint form, call

This institution is an equal opportunity provider.

A . de Conthe Rirst Year








Parenting can be hard work! Take care of yourself. It is easier to enjoy your baby and be a positive, loving parent when you are feeling good.



Exclusive breastfeeding recommended.

- Only breastmilk
- Mostly breastmilk with some iron-fortified formula
- o Iron-fortified formula
- No juice, water, cereal or food needed
- Ask doctor if extra vitamins or minerals are needed





Baby is ready to eat food.

- Breastfeed or formula feed first
- Use a small dish and feed by spoon only:
 - infant cereal
 - meat
 - fruits
 - vegetables
- Use single item baby foods or mashed table foods
- Try one new food at a time for 3-4 days
- Try food in any order
- Start water in a cup without a lid

when my baby is ready for new foods?

Begin trying finger foods



Continue to breastfeed or formula feed.

- Let baby use open cup for pumped breastmilk, formula, or water
- Let baby practice using a spoon
- Try soft table foods in small pieces, such as:
 - cooked vegetables
 - peeled, soft fruit
 - mashed or finely cut meat, chicken, fish, tofu, eggs
 - cheese slices, yogurt
 - toast
 - cottage cheese

DO NOT give hot dogs, peanut butter, nuts, seeds, popcorn, and other small, hard foods that can cause baby to choke.



- It is normal to continue breastfeeding past one year.
- Wean from bottle Child has outgrown the bottle
- Start whole milk in a cup
- Give table foods during family meals
- o Can offer small amounts of juice in a cup



- 1. Baby sits with support and has good head and neck control.
- 2. Baby leans forward with open mouth.
- 3. Baby can keep most of the food in his mouth and swallow it.
- 4. Baby can pick up food with his hand.